§ 180.554

Commodity	Parts per million
Cilantro, leaves	30.0
Cucumber	2.0
Fruit, stone, group 12, except plum, prune,	
fresh, postharvest	10.0
Ginseng	0.3
Grape	4.0
Grape, raisin	6.0
Juneberry	5.0
Kiwifruit, postharvest	15.0
Leafy greens subgroup 4A, except spinach	30.0
Lingonberry	5.0
Pear	10
Pepper, nonbell	0.02
Pistachio	0.02
Plum, prune, dried	2.5
Plum, prune, fresh	1.5
Pomegranate	2.0
Salal	5.0
Strawberry	3.0
Vegetable, fruiting, group 8, except nonbell	
pepper	2.0

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

[64 FR 28924, May 28, 1999, as amended at 65 FR 19849, Apr. 13, 2000; 65 FR 69883, Nov. 21, 2000; 67 FR 19120, Apr. 18, 2002; 68 FR 2247, Jan. 16, 2003; 68 FR 55519, Sept. 26, 2003; 71 FR 15617, Mar. 29, 2006; 71 FR 43664, Aug. 2, 2006; 73 FR 19154, Apr. 9, 2008]

§ 180.554 Kresoxim-methyl; tolerances for residues.

(a) General. (1) Tolerances are established for the combined residues of the fungicide kresoxim-methyl (methyl (E)-2-[2-(2-methylphenoxy)-methyl]phenyl-2-(methoxyimido)acetate) and its metabolites as follows: (E)-2-[2-(2-methylphenoxy)methyl]-phenyl-2-(methoxyimido)acetic acid; (E)-2-[2-(2-hydroxymethylphenoxy)methyl]-phenyl-2-(methoxyimido)acetic acid (free and glucose conjugated); and (E)-2-[2-(4-hydroxy-2-methylphenoxy)-methyl]phenyl-2-(methoxyimido)acetic acid (free and glucose conjugated) in or on the following commodities:

Commodity	Parts per million
Apple, dry pomace	1.0
Apple, wet pomace	1.0
Fruit, pome	0.5
Grape	1.0
Grape, raisin	1.5
Pecan	0.15
Vegetable, cucurbit, group 9	0.40

(2) Tolerances are established in or on the following commodities for the residues of the metabolite (*E*)-2-[2-(2-methylphenoxy)methyl]-phenyl-2-(methoxyimido)acetic acid resulting from the use of the fungicide kresoximmethyl:

Commodity	Parts per million
Cattle, meat byproducts Goat, meat byproducts Sheep, meat byproducts	0.01 0.01 0.01

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

[64 FR 31136, June 10, 1999, as amended at 71 FR 50359, Aug. 25, 2006; 74 FR 46377, Sept. 9, 2009]

§ 180.555 Trifloxystrobin; tolerances for residues.

(a) General. Tolerances are established for residues of trifloxystrobin, including its metabolites degradates, in or on the commodities in the table below. Compliance with the tolerance levels specified below is to be determined by measuring only the sum of trifloxystrobin, benzeneacetic acid, (E,E)- α -(methoxyimino)-2-[[[[1-[3phenyl]ethylidene] (trifluoromethyl) amino]oxy]methyl]-, methyl ester, and the free form of its acid metabolite CGA-321113, (E,E)-methoxyimino-[2-[1-

(3-trifluoromethyl-phenyl)ethylideneaminooxymethyl]phenyl]acetic acid, calculated as the stoichiometric equivalent of trifloxystrobin, in or on the com-

modity. Parts per Commodity Alfalfa, forage 0.01 Alfalfa, hay 0.01 Almond, hulls 9.0 Apple, wet pomace 5.0 Artichoke, globe 1.0 Asparagus 0.07 Banana 1 0.10 Barley, grain 0.05 Barley, hay 0.3 Barley, straw ...

0.4 0.2